AMENDMENTS TO THE CLAIMS:

The following is a complete list of the pending claims.

1. (Currently amended) A method for reducing the incorporation of non-standard amino acids into a heterologous protein expressed by microorganisms a microorganism

comprising:

- co-expressing in the microorganism at least one <u>a</u> heterologous protein and at least one <u>a</u> non-standard amino acid degrading protein.
- 2. (Currently amended) The method of claim 1 wherein the non-standard amino acid degrading protein is a glutamate dehydrogenase, leucine dehydrogenase, a valine dehydrogenase, a glutamate/leucine/phenylalanine/valine dehydrogenase, a phenylalanine dehydrogenase, or an opine dehydrogenase.
- 3. (Currently amended) The method of claim 2 wherein the non-standard amino acid degrading protein is a wild-type or K92L variant glutamate dehydrogenase from Escherichia coli, a leucine dehydrogenase from Bacillus cercus, a leucine dehydrogenase from Bacillus subtilis, a leucine dehydrogenase from Nostoc sp. a leucine dehydrogenase from Shewanella oneidensis, a valine dehydrogenase from Streptomyces avermitilis, or a glutamate/leucine/phenylalanine/valine dehydrogenase from Nitrosomonas curopaca.
- 4. (Currently amended) The method of claim 3 wherein the non-standard amino acid degrading protein has a sequence selected from SEQ ID NO:2[[,]] or 4, 6, 8, 10, 12, 14, or 16.
- 5. (Currently amended) The method of claim 4 wherein the non-standard amino acid degrading protein is encoded by a DNA molecule having a sequence selected from SEQ ID NO:1[[,]] or 3, 5, 7, 9, 11, 13, or 15.
- 6-7. (Cancelled)

- 8. (Original) The method of claim 1 wherein the microorganism is *Escherichia coli*.
- 9. (Currently amended) The method of claim 1 wherein at least one of the expressed heterologous protein(s) protein is a somatotropin.
- 10. (Currently amended) The method of claim 9 wherein the somatotropin is selected from the group consisting of human, equine, bovine, ovine, porcine, canine, [[or]] and feline somatotropin.
- 11. (Original) The method of claim 9 wherein the somatotropin is bovine somatotropin.
- 12. (Original) The method of claim 1 wherein the microorganism is *Escherichia coli* (*E. coli*); wherein the non-standard amino acid degrading protein is *E. coli* glutamate dehydrogenase or a lysine 92 leucine variant of *E. coli* glutamate dehydrogenase; and wherein the heterologous protein is bovine somatotropin.
- 13. (Withdrawn) The method of claim 1 wherein the non-standard amino acid degrading protein is a leucine dehydrogenase from *Bacillus cereus*, a leucine dehydrogenase from *Bacillus subtilis*, a leucine dehydrogenase from *Nostoc sp.*, a leucine dehydrogenase from *Shewanella oneidensis*, a valine dehydrogenase from *Streptomyces avermitilis*, or a glutamate/leucine/phenylalanine/valine dehydrogenase from *Nitrosomonas europaea*; and wherein the heterologous protein is bovine somatotropin.
- 14. (Original) The method of claim 1 wherein the heterologous protein and the non-standard amino acid degrading protein are expressed from a single expression vector.
- 15. (Original) The method of claim 1 wherein the heterologous protein and the non-standard amino acid degrading protein are expressed from at least two distinct expression vectors.

- 16. (Original) The method of claim 1 wherein the non-standard amino acid is norleucine.
- 17. (Currently amended) The method of claim 1 wherein the non-standard amino acid degrading protein is selected from the group consisting of: a glutamate dehydrogenase, a phenylalanine dehydrogenase, a valine dehydrogenase, a leucine dehydrogenase, a glutamate/leucine/phenylalanine/valine dehydrogenase and an opine dehydrogenase; and wherein the non-standard amino acid is selected from the group comprising: norleucine, norvaline, beta-methylnorleucine, and homoisoleucine.
- 18. (Original) The method of claim 17 wherein the non-standard amino acid is norleucine or norvaline.

19-26. (Cancelled)

- 27. (Currently amended) A method of isolating a protein from a microorganism comprising:
 - a) co-expressing in a microorganism at least one a heterologous protein and at least one
 a non-standard amino acid degrading protein; and
 - b) isolating the heterologous protein from the microorganism.
- 28. (Currently amended) A method of isolating a protein comprising:
 isolating at least one heterologous protein from a microorganism; wherein said
 microorganism co-expresses at least one <u>a</u> heterologous protein and at least one <u>a</u> nonstandard amino acid degrading protein.
- 29. (Currently amended) The method of either claim 27 or claim 28 wherein the non-standard amino acid degrading protein is selected from the group consisting of a glutamate dehydrogenase, a phenylalanine dehydrogenase, a leucine dehydrogenase, a valine dehydrogenase, a glutamate/leucine/phenylalanine/valine dehydrogenase, and an opine dehydrogenase.

- 30. (Original) The method of claim 29 wherein the non-standard amino acid is norleucine.
- 31. (Cancelled)
- 32. (Currently amended) The method of claim [[31]] 29 wherein the non-standard amino acid degrading enzyme protein is a wild-type or K92L variant glutamate dehydrogenase from Escherichia coli, a leucine dehydrogenase from Bacillus cercus, a leucine dehydrogenase from Bacillus subtilis, a leucine dehydrogenase from Nostoc sp. a leucine dehydrogenase from Shewanella oncidensis, a valine dehydrogenase from Streptomyces avermitilis, or a glutamate/leucine/phenylalanine/valine dehydrogenase from Nitrosomonas europaca.
- 33. (Cancelled)
- 34. (Withdrawn Currently amended) The method of claim [[31]] 27 wherein the non-standard amino acid degrading enzyme protein is a leucine dehydrogenase from Bacillus cereus, a leucine dehydrogenase from Bacillus subtilis, a leucine dehydrogenase from Nostoc sp. a leucine dehydrogenase from Shewanella oneidensis, a valine dehydrogenase from Streptomyces avermitilis, or a glutamate/leucine/phenylalanine/valine dehydrogenase from Nitrosomonas europaea.[[.]]
- 35. (Original) The method of either claim 27 or claim 28 wherein the microorganism is *Escherichia coli*.
- 36. (Currently amended) The method of claim 35 wherein the non-standard amino acid degrading enzyme protein is a glutamate dehydrogenase, a leucine dehydrogenase, a valine dehydrogenase, or a glutamate/leucine/phenylalanine/valine dehydrogenase.
- 37. (Currently amended) The method of either claim 27 or claim 28 wherein the microorganism is *Escherichia coli* (*E. coli*); and

wherein the non-standard amino acid degrading enzyme protein is a lysine 92 to leucine variant of *E. coli* glutamate dehydrogenase, a leucine dehydrogenase, a valine dehydrogenase, or glutamate/leucine/phenylalanine/valine dehydrogenase; and wherein at least one the heterologous protein is a bovine somatotropin.

- 38. (Currently amended) The method either of claims of either claim 27 or 28 wherein the non-standard amino acid degrading protein is selected from the group consisting of: a glutamate dehydrogenase, a phenylalanine dehydrogenase, a valine dehydrogenase, a leucine dehydrogenase, and a glutamate/leucine/phenylalanine/valine dehydrogenase; and wherein the non-standard amino acid is selected from the group comprising: norleucine, norvaline, beta-methylnorleucine, and homoisoleucine.
- 39. (Original) The method of claim 38 wherein the non-standard amino acid is norleucine or norvaline.
- 40. (Original) The method of either claim 27 or 28 wherein the non-standard amino acid degrading protein is capable of degrading norleucine.
- 41. (Original) The method of either claim 27 or 28 wherein the percentage of heterologous protein containing norleucine is substantially zero.
- 42. (Currently amended) The method of claim 1 wherein the heterologous protein and/or the norleucine non-standard amino acid degrading protein is expressed from a location in the microorganism's genome.
- 43. (Withdrawn New) The method of claim 1 wherein the non-standard amino acid degrading protein is a leucine dehydrogenase, a valine dehydrogenase, a glutamate/leucine/phenylalanine/valine dehydrogenase, a phenylalanine dehydrogenase, or an opine dehydrogenase.

- 44. (Withdrawn New) The method of claim 43 wherein the non-standard amino acid degrading protein is a leucine dehydrogenase from *Bacillus cereus*, a leucine dehydrogenase from *Bacillus subtilis*, a leucine dehydrogenase from *Nostoc sp.*, a leucine dehydrogenase from *Shewanella oneidensis*, a valine dehydrogenase from *Streptomyces avermitilis*, or a glutamate/leucine/phenylalanine/valine dehydrogenase from *Nitrosomonas europaea*.
- 45. (Withdrawn New) The method of claim 44 wherein the non-standard amino acid degrading protein has a sequence selected from SEQ ID NO:6, 8, 10, 12, 14, or 16.
- 46. (Withdrawn New) The method of claim 45 wherein the non-standard amino acid degrading protein is encoded by a DNA molecule having a sequence selected from SEQ ID NO:5, 7, 9, 11, 13, or 15.
- 47. (Withdrawn New) The method of claim 1 wherein the non-standard amino acid degrading protein is selected from the group consisting of: a phenylalanine dehydrogenase, a valine dehydrogenase, a leucine dehydrogenase, a glutamate/leucine/phenylalanine/valine dehydrogenase, and an opine dehydrogenase; and wherein the non-standard amino acid is selected from the group comprising: norleucine, norvaline, beta-methylnorleucine, and homoisoleucine.
- 48. (Withdrawn New) The method of claim 47 wherein the non-standard amino acid is norleucine or norvaline.